

THE FORMAMIDE₂-H₂O COMPLEX: STRUCTURE AND HYDROGEN BOND COOPERATIVE EFFECTS

SUSANA BLANCO

PABLO PINACHO

JUAN CARLOS LÓPEZ

Departamento de Química Física

Universidad de Valladolid, Valladolid

SPAIN



Universidad de Valladolid




Introduction

Peptide groups in biomolecules may adopt **cis** or **trans** configurations


These present a variety of hydrogen bond interactions sites:

- proton acceptor (**A**)
- proton donor (**D**)
- weak hydrogen bonds (**WD**) (if including the C_{α} -H groups)





Introduction

They made possible the interaction between peptide groups:
The $\text{C}=\text{O}\cdots\text{H}-\text{N}$ bond dominate in the hydrophobic core of proteins.




Introduction

These sites define a variety of possible interactions of peptide group with water: $\text{C=O}\cdots\text{HOH}$: ~44 %




Introduction

In proteins water may mediate in $\text{C}=\text{O}\cdots\text{H}-\text{N}$ to nucleate protein folding or in the protein surface where they are exposed to hydrophilic interactions



Introduction


Cooperative effects may play an important stabilization role:



G. A. Jeffrey "An Introduction to Hydrogen Bonding",
Oxford University Press, New York (1997).


Introduction

Cooperative effects may play an important stabilization role:




Introduction

Cooperative effects may play an important stabilization role:



Introduction

Cooperative effects may play an important stabilization role:



Resonance enhanced HB:

- Inductive polarization effects
- Shortens the N-C bond lengths
- Enlarges the C=O bond lengths

Previous studies

The interactions with water have been modelled from the study of gas phase **microsolvated** clusters of simple molecules having the peptide group:

Some examples from MW spectroscopy

formamide-(H₂O)_n, n=1,2 : F. J. Lovas et al. *J. Chem. Phys.* **1988**, 88, 722; S. Blanco *et al.*, *JACS, J. Am. Chem. Soc.* **2006**, 128, 12111.

N-methylformamide-(H₂O): W. Caminati, *et al.*, *Phys. Chem. Chem. Phys.* **2010**, 12, 10230–10234.

2-hydroxypyridine/2-pyridone-formamide-(H₂O)_n, n=1,2: A. Maris et al, *Chem. Phys. Lett.* 2002, 360, 155; S. Mata, *et al.* *J. Phys. Chem. A* 2010, 114, 11393–11398.



2-azetidinone-(H₂O)_n, n=1,2: J. C. López et al., *PCCP*, 2015, 17, 2054.

alaninamide-(H₂O): R. J. Lavrich, M. J. Tubergen, *J. Am. Chem. Soc.* 2000, 122, 2938–2943.


Formamide-water complexes

The observed **1:1** clusters of **formamide** with **water** cover all possible hydrogen bond situations:

trans-peptide




cis-peptide



42% in globular proteins

Formamide-water complexes

The signatures of polarization enhanced HB have observed when the structures of sequential 1:1 and 1:2 cycles:




Shortening of HB lengths

Stabilization energy

Formamide-water complexes

However no experimental evidence of changes in the $r(\text{C-N})$ or $r(\text{C=O})$ bond lengths.

Ab initio values



Formamide-water complexes


We have extended the study of formamide water clusters to new species:

(formamide)₂-H₂O¹ and formamide-(H₂O)₃

To characterize their structure and dynamics

To find the signatures of cooperative effects.

Spectrum



FTMW spectrometer

MB-FTMW (4-16 GHz)

Chem. Phys. 218 267 (1997); *Chem. Phys.* 119 880 (2003)

Summary

- Formamide₂-water (f₂w) can be taken as a microsolvated water cluster being formamide the solvent and as a model for water mediated amide interactions and self-association.
- Unambiguous structure characterization based on the observation of isotopologues and hyperfine structure analysis
- The structure of f₂w show for the first time in gas phase the effect of resonance enhanced cooperative hydrogen bonding of an amide
- Quadrupole coupling constants are a precious tool to detect the existence on inductive effects due to cooperativity polarization.

Acknowledgement

- Thank you for your attention!

